


that the sum of m plus n is 0, 1, 2 or 3; E, E<sup>I</sup>, E<sup>II</sup> and E<sup>III</sup> individually represent hydrogen or a suitable non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; and Q is selected from:



where Z' represents hydrogen or lower alkyl, acyl, alkoxy carbonyl, or aryloxy carbonyl; Z'' is hydrogen or lower alkyl; and Z''' is a non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; the dotted line indicates a carbon-carbon single bond or a carbon-carbon double bond, p is 0, 1 or 2; q is 0, 1, 2 or 3; and j is an integer from 0 to 3.

AI 2. (Amended) The compound of Claim 1 wherein X' is COR' where R' is selected from the group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic heterocyclylalkyl.

3. (Amended) The compound of Claim 2 wherein R' is phenyl or substituted phenyl.

A2 Sub B2 16. (Amended) A compound of the formula:

In re application of: DULL ET AL.


Serial No: 09/845,526

Filed: April 30, 2001

For: PHARMACEUTICAL COMPOSITIONS AND METHODS FOR USE

Examiner: V. Balasubramanian


Group Art Unit: 1624



where X'' is nitrogen, X and X' are individually carbon bonded to a substituent species selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, heterocyclyl, substituted heterocyclyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl; arylalkyl, substituted arylalkyl, halo, -OR', -NR'R'', -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, -C<sub>2</sub>R', -SR', -N<sub>3</sub>, C(=O)NR'R'', -NR'C(=O)R'', -C(=O)R', -C(=O)OR', -OC(=O)R', -O(CR'R''), C(=O)R', -O(CR'R''), NR'R'', -O(CR'R''), NR''C(=O)R', -O(CR'R''), NR''SO<sub>2</sub>R', -OC(=O)NR'R'', -NR'C(=O)O R'', -SO<sub>2</sub>R', -SO<sub>2</sub>NR'R'', and -NR'SO<sub>2</sub>R'', where R' and R'' are individually hydrogen, lower alkyl, cycloalkyl, heterocyclyl, or an aromatic group-containing species and r is an integer from 1 to 6, or R' and R'' can together form a cycloalkyl functionality; A is O, C=O or a covalent bond; D is a suitable non-hydrogen substituent species selected from the group of substituent species for X, X' and X''; k is 0, 1 or 2; Cx is selected from a group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic heterocyclylalkyl; m is an integer and n is an integer such that the sum of m plus n is 0, 1, 2 or 3; E, E', E'' and E''' individually represent hydrogen or a suitable non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; and Q is selected from:

In re application of: DULL ET AL.  
Serial No: 09/845,526  
Filed: April 30, 2001  
For: PHARMACEUTICAL COMPOSITIONS AND METHODS FOR USE

Examiner: V. Balasubramanian  
Group Art Unit: 1624



A3

22. (Amended) The compound of Claim 16 wherein X' is COR' where R' is selected from the group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic hetero-cyclylalkyl.


A4

24. (Amended) The compound of claim 16 selected from the group consisting of (S)-5-(2-pyrrolidin-2-ylethynyl)pyridine, (R)-5-(2-pyrrolidin-2-ylethynyl)pyridine, (S)-3-isopropoxy-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-phenyl-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-phenoxyphenyl-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-(4-methoxyphenyl)-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-(4-hydroxyphenyl)-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-cyclopentyloxy-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-cyclohexyloxy-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-(4-pyrrolidine-1-sulfonyl)phenoxy-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-pyridyloxy-5-(pyrrolidin-2-ylethynyl)pyridine, (S)-3-pyrrolidin-2-ylethynyl-5-(tetrahydropyran-4-yloxy)pyridine and (S)-3-(3,5-dihydroxy)phenoxy-5-(pyrrolidin-2-ylethynyl)pyridine.

In re application of: DULLE ET AL.  
 Serial No: 09/845,526  
 Filed: April 30, 2001  
 For: PHARMACEUTICAL COMPOSITIONS AND METHODS FOR USE

Examiner: V. Balasubramanian  
 Group Art Unit: 1624


25. (Amended) A pharmaceutical composition incorporating a compound of



where each of X and X' are individually carbon bonded to a substituent species selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, heterocyclyl, substituted heterocyclyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl; arylalkyl, substituted arylalkyl, halo, -OR', -NR'R'', -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, -C<sub>2</sub>R', -SR', -N<sub>3</sub>, C(=O)NR'R'', -NR'C(=O)R'', -C(=O)R', -C(=O)OR', -OC(=O)R', -O(CR'R'')<sub>r</sub>C(=O)R', -O(CR'R'')<sub>r</sub>NR'R'', -O(CR'R'')<sub>r</sub>NR'C(=O)R', -O(CR'R'')<sub>r</sub>NR'SO<sub>2</sub>R', -OC(=O)NR'R'', -NR'C(=O)OR'', -SO<sub>2</sub>R', -SO<sub>2</sub>NR'R'', and -NR'SO<sub>2</sub>R'', where R' and R'' are individually hydrogen, lower alkyl, cycloalkyl, heterocyclyl, or an aromatic group-containing species and r is an integer from 1 to 6, or R' and R'' can together form a cycloalkyl functionality; m is an integer and n is an integer such that the sum of m plus n is 0, 1, 2 or 3; E, E<sup>I</sup>, E<sup>II</sup> and E<sup>III</sup> individually represent hydrogen or a suitable non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; and Q is selected from:

In re application of: DULL ET AL.  
Serial No: 09/845,526  
Filed: April 30, 2001  
For: PHARMACEUTICAL COMPOSITIONS AND METHODS FOR USE

Examiner: V. Balasubramanian  
Group Art Unit: 1624



where Z' represents hydrogen or lower alkyl, acyl, alkoxycarbonyl, or aryloxycarbonyl; Z'' is hydrogen or lower alkyl; and Z''' is a non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; the dotted line indicates a carbon-carbon single bond or a carbon-carbon double bond; p is 0, 1 or 2; q is 0, 1, 2 or 3; and j is an integer from 0 to 3.

26. (Amended) The pharmaceutical composition of Claim 25 wherein X' is COR' where R' is selected from the group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic heterocyclylalkyl.

27. (Amended) The pharmaceutical composition of Claim 26 wherein R' is phenyl or substituted phenyl.

41. (Amended) A pharmaceutical composition incorporating a compound of the formula:

In re application of: DULL ET AL.


Serial No: 09/845,526

Filed: April 30, 2001

For: PHARMACEUTICAL COMPOSITIONS AND METHODS FOR USE


Examiner: V. Balasubramanian

Group Art Unit: 1624



AS


where X'' is nitrogen and X and X' are individually carbon bonded to a substituent species selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, heterocyclyl, substituted heterocyclyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl; arylalkyl, substituted arylalkyl, halo, -OR', -NR'R'', -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, -C<sub>2</sub>R', -SR', -N<sub>3</sub>, C(=O)NR'R'', -NR'C(=O)R'', -C(=O)R', -C(=O)OR', -OC(=O)R', -O(CR'R'')<sub>r</sub>C(=O)R', -O(CR'R'')<sub>r</sub>NR'R'', -O(CR'R'')<sub>r</sub>NR'C(=O)R', -O(CR'R'')<sub>r</sub>NR'SO<sub>2</sub>R', -OC(=O)NR'R'', -NR'C(=O)OR'', -SO<sub>2</sub>R', -SO<sub>2</sub>NR'R'', and -NR'SO<sub>2</sub>R'', where R' and R'' are individually hydrogen, lower alkyl, cycloalkyl, heterocyclyl, or an aromatic group-containing species and r is an integer from 1 to 6, or R' and R'' can together form a cycloalkyl functionality; A is O, C=O or a covalent bond; D is a suitable non-hydrogen substituent species selected from the group of substituent species for X, X' and X''; k is 0, 1 or 2; Cx is selected from a group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic hetero-cyclylalkyl; m is an integer and n is an integer such that the sum of m plus n is 0, 1, 2 or 3; E, E<sup>I</sup>, E<sup>II</sup> and E<sup>III</sup> individually represent hydrogen or a suitable non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; and Q is selected from:



where Z' represents hydrogen or lower alkyl, acyl, alkoxycarbonyl, or aryloxycarbonyl; Z'' is hydrogen or lower alkyl; and Z''' is a non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; the dotted line indicates a carbon-carbon single bond or a carbon-carbon double bond; p is 0, 1 or 2; q is 0, 1, 2 or 3; and j is an integer from 0 to 3.

A6 48. (Amended) The pharmaceutical composition of Claim 41 wherein X' is COR' where R' is selected from the group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic hetero-cyclylalkyl.


51. (Amended) A method for treating a central nervous system disorder, said method comprising administering an effective amount of a compound having the formula:



where each of X and X' are individually carbon bonded to a substituent species selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, heterocyclyl, substituted heterocyclyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, alkylaryl,

substituted alkylaryl; arylalkyl, substituted arylalkyl, halo, -OR', -NR'R'', -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, -C<sub>2</sub>R', -SR', -N<sub>3</sub>, C(=O)NR'R'', -NR'C(=O)R'', -C(=O)R', -C(=O)OR', -OC(=O)R', -O(CR'R'')<sub>r</sub>C(=O)R', -O(CR'R'')<sub>n</sub>NR'R'' -O(CR'R'')<sub>n</sub>NR'C(=O)R', -O(CR'R'')<sub>n</sub>NR'SO<sub>2</sub>R', -OC(=O)NR'R'', -NR'C(=O)OR'', -SO<sub>2</sub>R', -SO<sub>2</sub>NR'R'', and -NR'SO<sub>2</sub>R'', where R' and R'' are individually hydrogen, lower alkyl, cycloalkyl, heterocyclyl, or an aromatic group-containing species and r is an integer from 1 to 6, or R' and R'' can together form a cycloalkyl functionality; m is an integer and n is an integer such that the sum of m plus n is 0, 1, 2 or 3; E, E<sup>I</sup>, E<sup>II</sup> and E<sup>III</sup> individually represent hydrogen or a suitable non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; and Q is selected from:

Sub  
 135



where Z' is hydrogen, lower alkyl, acyl, alkoxy carbonyl, or aryloxy carbonyl; Z'' is hydrogen or lower alkyl; and Z''' is a non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; the dotted line indicates a carbon-carbon single bond or a carbon-carbon double bond; p is 0, 1 or 2; q is 0, 1, 2 or 3; and j is an integer from 0 to 3.

52. (Amended) The method of Claim 51 wherein X' is COR' where R' is selected from the group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic



In re application of: **DULL ET AL.**

Serial No: **09/845,526**

Filed: **April 30, 2001**

For: **PHARMACEUTICAL COMPOSITIONS AND METHODS FOR USE**


Examiner: **V. Balasubramanian**

Group Art Unit: **1624**

**A7** heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic heterocyclylalkyl.

53. (Amended) The method of Claim 51 wherein R' is phenyl or substituted phenyl.

66. (Amended) A method for treating a central nervous system disorder, said method comprising of the administration of an effective amount of a compound having the formula:



where X'' is nitrogen, X and X' are individually carbon bonded to a substituent species selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, heterocyclyl, substituted heterocyclyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl; arylalkyl, substituted arylalkyl, halo, -OR', -NR'R'', -CF<sub>3</sub>, -CN, -NO<sub>2</sub>, -C<sub>2</sub>R', -SR', -N<sub>3</sub>, C(=O)NR'R'', -NR'C(=O)R'', -C(=O)R', -C(=O)OR', -OC(=O)R', -O(CR'R'')<sub>r</sub>C(=O)R', -O(CR'R'')<sub>r</sub>NR'R'', -O(CR'R'')<sub>r</sub>NR''C(=O)R', -O(CR'R'')<sub>r</sub>NR''SO<sub>2</sub>R', -OC(=O)NR'R'', -NR'C(=O)OR'', -SO<sub>2</sub>R', -SO<sub>2</sub>NR'R'', and -NR'SO<sub>2</sub>R'', where R' and R'' are individually hydrogen, lower alkyl, cycloalkyl, heterocyclyl, or an aromatic group-containing species and r is an integer from 1 to 6, or R' and R'' can together form a cycloalkyl functionality; A is O, C=O or a covalent bond; D is a suitable non-hydrogen substituent species selected from the group of substituent species for X, X' and X''; k is 0, 1 or 2; Cx is selected from a group consisting of aryl, substituted aryl, heteroaryl, substituted heteroaryl, non-aromatic heterocyclyl, substituted non-aromatic heterocyclyl, non-aromatic heterocyclylalkyl and substituted non-aromatic hetero-cyclylalkyl; m is an integer and n is an integer such that the sum of m plus n is 0, 1, 2 or 3; E, E<sup>I</sup>, E<sup>II</sup> and E<sup>III</sup> individually represent hydrogen or a suitable non-hydrogen substituent selected from the group consisting of alkyl, substituted alkyl, halo-substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, alkylaryl, substituted alkylaryl, arylalkyl and substituted arylalkyl; and Q is selected from: